

Module 5: Schlieren and Shadowgraph

References

The Lecture Contains:

- References For Interferometry, Monochrome, Schlieren and Shadowgraph
- References For Color Schlieren

 **Previous**   **Next** 

**Module 5: Schlieren and Shadowgraph****References****FOR INTERFEROMETRY, MONOCHROME SCHLIEREN AND SHADOWGRAPH**

- G.S. Settles, *Schlieren and shadowgraph techniques: Visualizing phenomena in transport media* (Springer-Verlag publications, New York, 2001).
- R.J. Goldstein, ed., *Fluid Mechanics Measurements*, second edition, (Taylor and Francis, New York, 1996).
- F. Natterer, *The Mathematics of Computerized Tomography* (John Wiley, New York, 1986).
- G.T. Herman, *Image Reconstruction from Projections* (Academic, 1980).
- F. Mayinger, ed., *Optical Measurements: Techniques and Applications* (Springer-Verlag, Berlin, Germany, 1994).
- H. Lenski and M. Braun, Laser beam deflection: A method to investigate convection in vapor growth experiments, Proc. SPIE, 1557, Crystal Growth in Space & Related Optical Diagnostics, J. D. Trolinger & R. B. Lal (Eds.) (1991), 124-131.
- W. Schopf, J. C. Patterson and A. M. H. Brooker, Evaluation of the shadowgraph method for the convective flow in a side-heated cavity, Expts in Fluids, 21 (1996) 331-340.
- A dark field-type schlieren microscope for quantitative, in-situ mapping of solute concentration profiles around growing crystals, J. Cryst. Growth, 179 (1-2) (1997) 240-248.
- A. Srivastava, K. Muralidhar and P.K. Panigrahi, Comparison of interferometry, schlieren and shadowgraph for visualizing convection around a KDP crystal, J. Cryst. Growth, 267 (2004) 348-361.
- Masayuki Mantani, Motoyuki Sugiyama and Tomoya Ogawa, Electronic measurement of concentration gradient around a crystal growing from a solution by using Mach-Zehnder interferometer, J. Cryst. Growth, 114 (1991) 71-76.

A set of small navigation icons consisting of arrows pointing left and right, used for navigating between pages or sections.

**Module 5: Schlieren and Shadowgraph****References**

- Y. Suzuki, T. Sawada, S. Miyashita and H. Komatsu, In-situ measurements of the solubility of crystals under high pressure by an interferometric method, Rev. Sci. Instruments, 69 (7) (1998) 2720-2724.
- T. Inoue, K. Mori, Y. Kageyama, and H. Mori, Various growth shapes of Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub>.5H<sub>2</sub>O crystals, Cryst. Res. Technol., 35 (5) (2000) 587-593.
- K. Nagashima and Y. Furukawa, Time development of a solute diffusion field and morphological instability on a planar interface in the directional growth of ice crystals, J. Cryst. Growth, 209 (2000) 167-174.
- K. Onuma, K. Tsukamoto and S. Nakadate, Application of real time phase shift interferometer to the measurement of concentration field, J. Cryst. Growth, 129 (1993) 706-718.
- L. Duan and J. Z. Shu, The convection during NaClO<sub>3</sub> crystal growth observed by th K. Onuma, T. Kameyama and K. Tsukamoto, In-situ study of surface phenomena by real time phase shift interferometry, JCG, 137 (1994) 610-622
- H. Komatsu, S. Miyashita and Y. Suzuki, Interferometric observation of the interfacial concentration gradient layers around a lysozyme crystal, Jpn. J. Appl. Phys., Part 2, 32 (12B) (1993) L1855-L1857.
- Gen Sasaki, Kazuo Kurihara, Toshitaka Nakada, Satoru Miyashita and Hiroshi Komatsu, A novel approach to the solubility measurement of protein crystals by two-beam interferometry, J. Cryst. Growth, 169 (1996) 355-360.
- G.W. Faris and R.L. Byer, Three-dimensional beam-deflection optical tomography of a supersonic jet, Appl. Opt., 27 (24) (1988) 5202-5212.
- Atul Srivastava, K. Muralidhar and P. K. Panigrahi, Reconstruction of the concentration field around a growing KDP crystal with schlieren tomography, Appl. Opt., 44 (26) (2005) 5381-5392.

A set of navigation icons consisting of arrows pointing left and right, with the word "Previous" and "Next" centered between them.

**Module 5: Schlieren and Shadowgraph****References****FOR COLOR SCHLIEREN**

- M. Mantani, M. Sugiyama, T. Ogawa, Electronic measurement of concentration gradient around a crystal growing from a solution by using Mach-Zehnder interferometer, Journal of Crystal Growth, Vol. 114 (1991) p. 71.
- K. Onuma, T. Tsukamoto, S. Nakadate, Application of real time phase shift interferometer to the measurement of concentration field, Journal of Crystal Growth, Vol. 129 (1993) p. 706.
- A.K. Shenoy, A.K. Agrawal, S.R. Gollahalli, Quantitative evaluation of flow computations by rainbow schlieren deflectometry, AIAA Journal, Vol. 36 (1998) p. 1953.
- G. Tanda, F. Devia, Application of a schlieren technique to heat transfer measurements in free convection, Experiments in Fluids, Vol. 24 (1998) p. 285.
- A.M. Schwartz, K.A. Berglund, The use of Raman spectroscopy for in situ monitoring of lysozyme concentration during crystallization in a hanging drop, Journal of Crystal Growth, Vol. 203 (1999) p. 599.
- V.I. Bredikhin, G.L. Galushkina, S.P. Kuznetsov, Schlieren technique to in situ monitor-rapidly growing KDP crystal surface, Journal of Crystal Growth, Vol. 219 (2000) p. 83.
- A. Srivastava, A. Phukan, P.K. Panigrahi, K. Muralidhar, Imaging of a convective field in a rectangular cavity using interferometry, schlieren and shadowgraph, Optics and Lasers in Engineering, Vol. 42 (2004) p. 469.
- P.S. Greenberg, R.B. Klimek, R. Buchele, Quantitative rainbow schlieren deflectometry, Applied Optics, Vol. 34 (19) (1995) p. 3810.
- N.P. Zaitseva, L.N. Ranckovich, S.V. Bogatyreva, Stability of KH<sub>2</sub>PO<sub>4</sub> and K(HD)<sub>2</sub>PO<sub>4</sub> solutions at fast crystal growth rates, Journal of Crystal Growth Vol. 148 (1995) p. 276.

Previous    Next