

Discrete Mathematics

Lecture 12: Proof Technique (Contrapositive)

Instructor: Sourav Chakraborty

Proof Techniques

To prove statement $A \implies B$.

There are different proof techniques:

- Constructive Proofs
- Proof by Contradiction
- Proof by Contrapositive
- Induction
- Counter example
- Existential Proof

Which approach to apply

Which approach to apply

- It depends on the problem.

Which approach to apply

- It depends on the problem.
- Sometimes the problem can be split into smaller problems that can be easier to tackle individually.

Which approach to apply

- It depends on the problem.
- Sometimes the problem can be split into smaller problems that can be easier to tackle individually.
- Sometimes viewing the problem in a different way can also help in tackling the problem easily.

Which approach to apply

- It depends on the problem.
- Sometimes the problem can be split into smaller problems that can be easier to tackle individually.
- Sometimes viewing the problem in a different way can also help in tackling the problem easily.
- Whether to split a problem or how to split a problem or how to look at a problem is an ART that has to be developed.

Which approach to apply

- It depends on the problem.
- Sometimes the problem can be split into smaller problems that can be easier to tackle individually.
- Sometimes viewing the problem in a different way can also help in tackling the problem easily.
- Whether to split a problem or how to split a problem or how to look at a problem is an ART that has to be developed.
- There are some thumb rules but at the end it is a skill you develop using a lot of practice.

Tricks for solving problems

- (Splitting into smaller problem) If the problem is to prove $A \implies B$ and B can be written as $B = C \wedge D$ then note that

$$(A \implies B) \equiv (A \implies C \wedge D) \equiv (A \implies C) \wedge (A \implies D).$$

- (Remove Redundant Assumptions) If $A \implies B$ then $A \wedge C$ also implies B .

Tricks for solving problems

- (Splitting into smaller problem) If the problem is to prove $A \implies B$ and B can be written as $B = C \wedge D$ then note that

$$(A \implies B) \equiv (A \implies C \wedge D) \equiv (A \implies C) \wedge (A \implies D).$$

- (Remove Redundant Assumptions) If $A \implies B$ then $A \wedge C$ also implies B .

$$(A \implies B) \implies (A \wedge C \implies B) = \text{True}$$

Tricks for solving problems

- (Splitting into smaller problem) If the problem is to prove $A \implies B$ and B can be written as $B = C \wedge D$ then note that

$$(A \implies B) \equiv (A \implies C \wedge D) \equiv (A \implies C) \wedge (A \implies D).$$

- (Remove Redundant Assumptions) If $A \implies B$ then $A \wedge C$ also implies B .

$$(A \implies B) \implies (A \wedge C \implies B) = \text{True}$$

- (Sometimes proving something stronger is easier) If $C \implies B$ then

$$(A \implies C) \implies (A \implies B).$$

Constructive Proof: Direct Proof

- For proving $A \implies B$ we can start with the assumption A and step-by-step prove that B is true.

Constructive Proof: Direct Proof

- For proving $A \implies B$ we can start with the assumption A and step-by-step prove that B is true.
- Sometimes a direct proof (as in the previous example) can be magical and hard to understand how to obtain.

Constructive Proof: Direct Proof

- For proving $A \implies B$ we can start with the assumption A and step-by-step prove that B is true.
- Sometimes a direct proof (as in the previous example) can be magical and hard to understand how to obtain.
- A simpler technique is to have a backward proof.

Constructive Proof: Direct Proof

- For proving $A \implies B$ we can start with the assumption A and step-by-step prove that B is true.
- Sometimes a direct proof (as in the previous example) can be magical and hard to understand how to obtain.
- A simpler technique is to have a backward proof.
- If we have to prove $(A \implies B)$ then the idea is to simplify B .

Constructive Proof: Direct Proof

- For proving $A \implies B$ we can start with the assumption A and step-by-step prove that B is true.
- Sometimes a direct proof (as in the previous example) can be magical and hard to understand how to obtain.
- A simpler technique is to have a backward proof.
- If we have to prove $(A \implies B)$ then the idea is to simplify B .
- And if $C \iff B$ then $(A \implies B) \equiv (A \implies C)$.

Constructive Proof: Case Studies

Constructive Proof: Case Studies

- Sometimes the assumption or the premise can be split into different cases. In that case we can split the problem according to cases.

Constructive Proof: Case Studies

- Sometimes the assumption or the premise can be split into different cases. In that case we can split the problem according to cases.
- If $A = C \vee D$ then

$$(A \implies B) \equiv (C \implies B) \wedge (D \implies B).$$

Proof by Contradiction

- Note that

$$(A \implies B) \equiv (\neg B \wedge A = \text{False})$$

This is called “proof by contradiction”

- To proof $A \implies B$ sometimes its easier to prove that

$$\neg B \wedge A = \text{False}.$$

- A similar statement is

$$(A \implies B) \equiv (\neg B \implies \neg A)$$

This is called “proof by contra-positive”

Contra-positive Proof

- A similar statement is

$$(A \implies B) \equiv (\neg B \implies \neg A)$$

This is called “proof by contra-positive”

Contra-positive Proof

- A similar statement is

$$(A \implies B) \equiv (\neg B \implies \neg A)$$

This is called “**proof by contra-positive**”

- This is particularly useful when B (the deduction) is of the form $C \vee D$

Contra-positive Proof

- A similar statement is

$$(A \implies B) \equiv (\neg B \implies \neg A)$$

This is called “**proof by contra-positive**”

- This is particularly useful when B (the deduction) is of the form $C \vee D$
- In that case

$$(A \implies B) \equiv (\neg B \implies \neg A) \equiv ((\neg C \wedge \neg D) \implies \neg A)$$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

- $A = “a^2 + b^2 \text{ is even}”$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

- $A = “a^2 + b^2 \text{ is even}”$
- $B = “\text{either both } a \text{ and } b \text{ are odd or both } a \text{ and } b \text{ are even.}”$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

- $A = “a^2 + b^2$ is even”
- $B = “\text{either both } a \text{ and } b \text{ are odd or both } a \text{ and } b \text{ are even.}”$
- $C = “\text{both } a \text{ and } b \text{ are odd}”$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

- $A = “a^2 + b^2$ is even”
- $B = “\text{either both } a \text{ and } b \text{ are odd or both } a \text{ and } b \text{ are even.}”$
- $C = “\text{both } a \text{ and } b \text{ are odd}”$
- $D = “\text{both } a \text{ and } b \text{ are even}”$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

- $A = “a^2 + b^2 \text{ is even}”$
- $B = “\text{either both } a \text{ and } b \text{ are odd or both } a \text{ and } b \text{ are even.}”$
- $C = “\text{both } a \text{ and } b \text{ are odd}”$
- $D = “\text{both } a \text{ and } b \text{ are even}”$

$$(A \implies B) \equiv (\neg B \implies \neg A) \equiv ((\neg C \wedge \neg D) \implies \neg A)$$

Example 1

Problem

If a and b are two positive integers and $a^2 + b^2$ is even then either both a and b are odd or both a and b are even.

Is same as ...

Problem

If a and b are two positive integers and one of a or b is odd and the other is even then $a^2 + b^2$ cannot be even.

Example 1

Problem

If a and b are two positive integers and one of a or b is odd and the other is even then $a^2 + b^2$ cannot be even.

Proof:

Example 1

Problem

If a and b are two positive integers and one of a or b is odd and the other is even then $a^2 + b^2$ cannot be even.

Proof:

Proof using case studies. There are two cases:

Example 1

Problem

If a and b are two positive integers and one of a or b is odd and the other is even then $a^2 + b^2$ cannot be even.

Proof:

Proof using case studies. There are two cases:

- 1 a is odd and b is even
- 2 a is even and b is odd.

Example 1

Problem

If a and b are two positive integers and one of a or b is odd and the other is even then $a^2 + b^2$ cannot be even.

Proof:

Proof using case studies. There are two cases:

- 1 a is odd and b is even
- 2 a is even and b is odd.

Complete the proof by yourself.

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

- $A =$ “the product ab is an irrational number”

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

- A = “the product ab is an irrational number”
- B = “either a or b must be an irrational number.”

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

- A = “the product ab is an irrational number”
- B = “either a or b must be an irrational number.”
- C = “ a must be an irrational number”

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

- A = “the product ab is an irrational number”
- B = “either a or b must be an irrational number.”
- C = “ a must be an irrational number”
- D = “ b must be an irrational number”

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

- A = “the product ab is an irrational number”
- B = “either a or b must be an irrational number.”
- C = “ a must be an irrational number”
- D = “ b must be an irrational number”

$$(A \implies B) \equiv (\neg B \implies \neg A) \equiv ((\neg C \wedge \neg D) \implies \neg A)$$

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

This is same as...

Example 2

Problem

If a and b are real numbers such that the product ab is an irrational number, then either a or b must be an irrational number.

This is same as...

Problem

If a and b are rational numbers then ab is a rational number.

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

- $A = “n \equiv 2 \pmod{3}”$

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

- $A = “n \equiv 2 \pmod{3}”$
- $B = “n$ is not a square of an integer.”

Example 3

Problem

If n is a positive integer such that $n \equiv 2(\text{mod } 3)$, then n is not a square of an integer.

- $A = “n \equiv 2(\text{mod } 3)”$
- $B = “n$ is not a square of an integer.”

$$(A \implies B) \equiv (\neg B \implies \neg A) \equiv ((\neg C \wedge \neg D) \implies \neg A)$$

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

This is same as ...

Example 3

Problem

If n is a positive integer such that $n \equiv 2 \pmod{3}$, then n is not a square of an integer.

This is same as ...

Problem

If n is a positive integer and n is a square of an integer then $n \not\equiv 2 \pmod{3}$.

Example 3

Problem

If n is a positive integer and n is a square of an integer then $n \not\equiv 2 \pmod{3}$.

Example 3

Problem

If n is a positive integer and n is a square of an integer then $n \not\equiv 2 \pmod{3}$.

How do you solve this problem?

Example 3

Problem

If n is a positive integer and n is a square of an integer then $n \not\equiv 2 \pmod{3}$.

How do you solve this problem?

This is an exercise. We will discuss it in the problem solving video.