

Discrete Mathematics

Lecture 4: Propositional Logic and Predicate Logic

Instructor: Sourav Chakraborty

Propositional logic and Predicate Logic

Propositional logic and Predicate Logic

- Every statement (or proposition) is either TRUE or FALSE.

Propositional logic and Predicate Logic

- Every statement (or proposition) is either TRUE or FALSE.
- A statement can be formed using other statements connected to each other by 5 kinds of connectives: AND, OR, NOT, IMPLIES and IFF.

Propositional logic and Predicate Logic

- Every statement (or proposition) is either TRUE or FALSE.
- A statement can be formed using other statements connected to each other by 5 kinds of connectives: AND, OR, NOT, IMPLIES and IFF.
- A statement can have unspecified terms, called variables. All the variables has to be properly quantified using the two quantifiers FOR-ALL and THERE EXISTS.

Checking correctness of a statement or theorem

- Any meaningful statement / proposition / theorem can be written as a mathematical logic statement.

Checking correctness of a statement or theorem

- Any meaningful statement / proposition / theorem can be written as a mathematical logic statement.
- A statement is consistent or correct if for any setting of the input variables (smaller statements) to TRUE or FALSE the statement always evaluate to TRUE.

Checking correctness of a statement or theorem

- Any meaningful statement / proposition / theorem can be written as a mathematical logic statement.
- A statement is consistent or correct if for any setting of the input variables (smaller statements) to TRUE or FALSE the statement always evaluate to TRUE.

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Is $A \implies B$ is equivalent to $(\neg B \wedge A) = FALSE$

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Is $A \implies B$ is equivalent to $(\neg B \wedge A) = FALSE$

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Is $A \implies B$ is equivalent to $(\neg B \wedge A) = FALSE$

A	B	$A \implies B$	$(\neg B \wedge A) = F$
F	F		
F	T		
T	F		
T	T		

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Is $A \implies B$ is equivalent to $(\neg B \wedge A) = FALSE$

A	B	$A \implies B$	$(\neg B \wedge A) = F$
F	F	T	
F	T	T	
T	F	F	
T	T	T	

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Is $A \implies B$ is equivalent to $(\neg B \wedge A) = FALSE$

A	B	$A \implies B$	$(\neg B \wedge A) = F$
F	F	T	T
F	T	T	T
T	F	F	F
T	T	T	T

Problems on Propositional Logic

- Prove that $(a \implies b)$ is equivalent to $(\neg a \vee b)$.
- Prove that $(a \iff b)$ is equivalent to $(a \implies b) \wedge (b \implies a)$.
- Prove that $(p \vee q) \implies r$ is equivalent to $(p \implies r) \wedge (q \implies r)$.
- Prove that $(a \implies b)$ is equivalent to $(\neg b \implies \neg a)$.

Checking Equivalence

Definition

Two statements are equivalent if their TRUTH TABLES are the same.

Checking Equivalence

Definition

Two statements are equivalent if their TRUTHTABLES are the same.

Another approach is to use some already proved rules to simplify the formulas before using the brute force truth table approach.

Rules of Propositional Logic

Let p , q and r be propositions.

- ❶ Commutative law:

$$(p \vee q) = (q \vee p) \text{ and } (p \wedge q) = (q \wedge p)$$

- ❷ Associative law:

$$(p \vee (q \vee r)) = ((p \vee q) \vee r) \text{ and } (p \wedge (q \wedge r)) = ((p \wedge q) \wedge r)$$

- ❸ Distributive law:

$$(p \vee (q \wedge r)) = (p \vee q) \wedge (p \vee r) \text{ and} \\ (p \wedge (q \vee r)) = (p \wedge q) \vee (p \wedge r)$$

- ❹ De Morgan's Law:

$$\neg(p \vee q) = (\neg p \wedge \neg q) \text{ and } \neg(p \wedge q) = (\neg p \vee \neg q)$$

Rules for Negation

- $\neg(\forall x P(x))$ is same as $\exists x (\neg P(x))$
- $\neg(\exists x P(x))$ is same as $\forall x (\neg P(x))$

Problems on Propositional Logic

The function

$$\left((p \vee (r \vee q)) \wedge \neg(p \wedge (\neg q \wedge \neg r)) \right)$$

is equal to which of the following functions:

- A. $q \vee r$
- B. $\neg p \vee (r \wedge q)$
- C. $(p \vee q) \vee r$
- D. $(p \vee q) \wedge \neg(p \vee r)$
- E. $(p \wedge r) \vee (p \wedge q)$

Problems on Propositional Logic

The function

$$\left((p \vee (r \vee q)) \wedge \neg(p \wedge (\neg q \wedge \neg r)) \right)$$

Problems on Propositional Logic

The function

$$\left((p \vee (r \vee q)) \wedge \neg(p \wedge (\neg q \wedge \neg r)) \right)$$

Problems on Propositional Logic

The function

$$\left((p \vee (r \vee q)) \wedge \neg(p \wedge (\neg q \wedge \neg r)) \right)$$

is equal to which of the following functions:

- A. $q \vee r$
- B. $\neg p \vee (r \wedge q)$
- C. $(p \vee q) \vee r$
- D. $(p \vee q) \wedge \neg(p \vee r)$
- E. $(p \wedge r) \vee (p \wedge q)$

Negating a sentence

What is the negation of the sentence: “There is an university in USA where every department has at least 20 faculty and at least one noble laureate.”

Negating a sentence

What is the negation of the sentence: “There is an university in USA where every department has at least 20 faculty and at least one noble laureate.”

- A. There is an university in USA where every department has less than 20 faculty and at least one noble laureate.
- B. All universities in USA where every department has at least 20 faculty and at least one noble laureate.
- C. For all universities in USA there is a department has less than 20 faculty or at most one noble laureate.
- D. For all universities in USA there is a department has less than 20 faculty and at least one noble laureate.

Propositional Logic and Predicate Logic

- Every statement is either TRUE or FALSE
- There are logical connectives \vee , \wedge , \neg , \implies and \iff .
- Two logical statements can be equivalent if the two statements answer exactly in the same way on every input.
- To check whether two logical statements are equivalent one can do one of the following:
 - Checking the Truthtable of each statement
 - Reducing one to the other using reductions

Propositional Logic and Predicate Logic

- There are two important symbols: \forall and \exists .
- Some statements can be defined using a variable.
- For example: $P_x = "4x^2 + 3 \text{ is divisible by } 5"$
- We can have statements like: $\forall x \in \mathbb{Z}, 4x^2 + 3 \text{ is divisible by } 5$.
- Or $\exists x \in \mathbb{Z}, 4x^2 + 3 \text{ is divisible by } 5$.

Using Propositional Logic for designing proofs

- A mathematical statement comprises of a premise (or assumptions). And when the assumptions are satisfied the statement deduces something.

Using Propositional Logic for designing proofs

- A mathematical statement comprises of a premise (or assumptions). And when the assumptions are satisfied the statement deduces something.
- If A is the set of assumptions and B is the deduction then a mathematical statement is of the form

$$A \implies B$$

Using Propositional Logic for designing proofs

- A mathematical statement comprises of a premise (or assumptions). And when the assumptions are satisfied the statement deduces something.
- If A is the set of assumptions and B is the deduction then a mathematical statement is of the form

$$A \implies B$$

- Now how to check if the statement is correct? And if it is indeed correct how to prove the statement?

Using Propositional Logic for designing proofs

- A mathematical statement comprises of a premise (or assumptions). And when the assumptions are satisfied the statement deduces something.
- If A is the set of assumptions and B is the deduction then a mathematical statement is of the form

$$A \implies B$$

- Now how to check if the statement is correct? And if it is indeed correct how to prove the statement?
- Depending on whether A or B (or both) can be split into smaller statements and how the smaller statements are connected we can design different techniques for proving the overall statement of $A \implies B$.

Using Propositional Logic for designing proofs

- A mathematical statement comprises of a premise (or assumptions). And when the assumptions are satisfied the statement deduces something.
- If A is the set of assumptions and B is the deduction then a mathematical statement is of the form

$$A \implies B$$

- Now how to check if the statement is correct? And if it is indeed correct how to prove the statement?
- Depending on whether A or B (or both) can be split into smaller statements and how the smaller statements are connected we can design different techniques for proving the overall statement of $A \implies B$.
- If indeed we can prove that the statement is correct then we can call it a Theorem.

Proof Techniques

To prove statement B from A .

- Constructive Proofs
- Proof by Contradiction
- Proof by Contrapositive
- Induction
- Counter example
- Existential Proof